

Stat 8112 Lecture Notes  
**Big Oh Pee and Little Oh Pee**  
Charles J. Geyer  
January 23, 2013

## 1 Big Oh and Little Oh

A sequence  $x_n$  of non-random vectors is said to be  $O(1)$  if it is bounded and  $o(1)$  if it converges to zero. If  $a_n$  is a sequence of non-random positive scalars, then

$$x_n = O(a_n) \tag{1a}$$

means

$$\frac{x_n}{a_n} = O(1)$$

(that is,  $x_n/a_n$  is bounded), and

$$x_n = o(a_n) \tag{1b}$$

means

$$\frac{x_n}{a_n} = o(1)$$

(that is,  $x_n/a_n$  converges to zero).

Note that the equals sign in (1a) and (1b) don't actually mean that anything is equal to anything. These "equations" are a convenient shorthand, but they are not about equality. Thus in careful arguments one needs to replace  $O(a_n)$  and  $o(a_n)$  by their definitions. Replace (1a) by there exists  $M < \infty$  such that

$$\|x_n\| \leq Ma_n, \quad \forall n \in \mathbb{N}.$$

Replace (1b) by

$$\frac{x_n}{a_n} \rightarrow 0, \quad n \rightarrow \infty.$$

It is sometimes convenient to allow the sequences  $a_n$  to be vector-valued, in which case  $O(a_n)$  means the same thing as  $O(\|a_n\|)$  and  $o(a_n)$  means the same thing as  $o(\|a_n\|)$ . That is, this is merely an abbreviation allowing us to omit the norm symbol.

We use the same notation to describe functions as well as sequences. We say that a function is  $O(1)$  as  $x \rightarrow 0$  if it is bounded on a neighborhood of zero, and we say it is  $o(1)$  as  $x \rightarrow 0$  if

$$f(x) \rightarrow 0, \quad \text{as } x \rightarrow 0.$$

For any other function  $g$

$$f(x) = O(g(x))$$

means

$$\frac{f(x)}{\|g(x)\|} = O(1),$$

and

$$f(x) = o(g(x))$$

means

$$\frac{f(x)}{\|g(x)\|} = o(1).$$

And we use similar notation for behavior at other points. For example, we say a Cauchy probability density function is  $O(x^{-2})$  as  $|x| \rightarrow \infty$ .

## 2 Big Oh Pee and Little Oh Pee

A sequence  $X_n$  of random vectors is said to be  $O_p(1)$  if it is bounded in probability (tight) and  $o_p(1)$  if it converges in probability to zero.

The notations gain power when we consider pairs of sequences. Suppose  $X_n$  and  $Y_n$  are random sequences taking values in any normed vector space, then

$$X_n = O_p(Y_n) \tag{2a}$$

means  $X_n/\|Y_n\|$  is bounded in probability and

$$X_n = o_p(Y_n) \tag{2b}$$

means  $X_n/\|Y_n\|$  converges in probability to zero.

These notations are often used when the sequence  $Y_n$  is deterministic, for example,  $X_n = O_p(n^{-1/2})$ . They are also often used when the sequence  $Y_n$  is random, for example, we say two estimators  $\hat{\theta}_n$  and  $\tilde{\theta}_n$  of a parameter  $\theta$  are *asymptotically equivalent* if

$$\hat{\theta}_n - \tilde{\theta}_n = o_p(\hat{\theta}_n - \theta)$$

and

$$\hat{\theta}_n - \tilde{\theta}_n = o_p(\tilde{\theta}_n - \theta).$$

Again the equals sign in (2a) and (2b) don't actually mean that anything is equal to anything. Thus in careful arguments one needs to replace (2a) by for every  $\varepsilon > 0$  there exists an  $M < \infty$  such that

$$\Pr(\|X_n\| \leq M\|Y_n\|) \geq 1 - \varepsilon, \quad \forall n \in \mathbb{N}$$

and replace (2b) by

$$\frac{X_n}{\|Y_n\|} \xrightarrow{P} 0, \quad n \rightarrow \infty$$

or by for every  $\varepsilon > 0$

$$\Pr(\|X_n\| \geq \varepsilon \|Y_n\|) \rightarrow 0.$$

We also use big oh and little oh and big oh pee and little oh pee notation for terms in equations (not just for right-hand sides of equations). For example, a function  $f$  is differentiable at  $x$  if

$$f(x+h) = f(x) + f'(x)h + o(h)$$

(see the handout on the delta method for more on this) and one case of Slutsky's theorem says

$$X_n \xrightarrow{w} X$$

implies

$$X_n + o_p(1) \xrightarrow{w} X.$$